

Gout management plan

Name: _____

GP: _____

Phone: _____

What causes gout?

Gout is a type of arthritis that is caused by too much uric acid building up in the body.

It is normal for the body to produce uric acid and pass it out in urine.

Uric acid may build up if you eat certain foods, take certain medications, or have kidney problems.

Allopurinol lowers uric acid. If you can keep your uric acid below 0.36 mmol/l it will help to stop your gout attacks.

This may take six to twelve months of taking allopurinol every day.

Help yourself to prevent gout attacks

Eat regular meals

- Starving or feasting may cause gout attacks

Avoid or limit food that can cause gout

- Red meat, shellfish, offal

If you are overweight, losing weight can help to lower uric acid

- Eat smaller meals
- 30 minutes of physical activity daily

Limit alcohol and sugary drinks

- This includes fruit juice

Drink plenty of water or trim milk

- To help flush uric acid out of your body

Produced by the Department of Rheumatology, Immunology and Allergy, Christchurch Hospital, Private Bag 4710, Christchurch, New Zealand

Take allopurinol every day

Allopurinol tablets come in two strengths: 100 mg tablets and 300 mg tablets. Start at a low dose and build up as prescribed:

Weeks 1 – 4 _____ mg/day

Weeks 5 – 8 _____ mg/day

Weeks 9 – _____ mg/day

Continue on this dose.

Allopurinol can trigger attacks of gout when you first start taking it.

If you get a rash or an itch, STOP allopurinol immediately and see your doctor.

Target uric acid: 0.36 mmol/l or less

Your uric acid:	Date:
_____ mmol/l	____/____/____
_____ mmol/l	____/____/____
_____ mmol/l	____/____/____
_____ mmol/l	____/____/____
_____ mmol/l	____/____/____
_____ mmol/l	____/____/____
_____ mmol/l	____/____/____
_____ mmol/l	____/____/____
_____ mmol/l	____/____/____

Acute gout attacks

What to do in an acute attack of gout?

Keep taking your allopurinol! DON'T STOP.

- Rest and use ice packs
- See your doctor if no better after 24 hours

You can take other medication to relieve the pain of acute attacks or to help prevent them from happening.

Your reliever medication is: _____

Dose: _____

Take: _____